

What drives the weather changes

Gregory Falkovich

Dept of Physics, Weizmann Institute of Science

October 24, 2015 Kyiv

“The answer is blowing in the wind”

Hydrostatics

Only normal forces

$$\nabla p = \rho \mathbf{f}$$

$$\mathbf{f} = -\nabla \phi$$

$$\nabla \rho \times \nabla \phi = 0$$

$$\phi = gz$$

$$\partial p / \partial z = -\rho g$$

Horizontal temperature
gradient causes wind

$$T(z) = T_0 - \alpha z$$


$$p(z) = p(0)(1 - \alpha z / T_0)^{mg/\alpha}$$

$$\alpha \simeq 6.5^\circ / km$$


$$\rho(p', s) > \rho(p', s') \Rightarrow \left(\frac{\partial \rho}{\partial s} \right)_p \frac{ds}{dz} < 0$$

$$c_p dT < g dz$$

$$-\frac{dT}{dz} < \frac{g}{c_p} \sim 10^\circ / km$$



Horizontal temperature
gradient causes wind



$$\nabla \rho \times \nabla \phi \neq 0$$

4 25 2005 15307



24


8 km

NOAA

[HTTP://WWW.NOAA.GOV](http://www.noaa.gov)



Atmospheric spectrum



Nastrom, Gage,
J. Atmosph. Sci. 1985

Atmospheric flows are driven by the gradients of solar heating.

Vertical gradients cause thermal convection on the scale of the troposphere depth (less than 10 km).


Horizontal gradients excite motions on a planetary (10000 km) and smaller scales.

Weather is mostly determined by the flows at intermediate scale (hundreds of kilometers).


Where these flows get their energy from?

The puzzle is that three-dimensional small-scale motions cannot transfer energy to larger scales while large-scale planar motions cannot transfer energy to smaller scales.



Three- dimensional turbulence: fragmentation and energy transfer to smaller scales




Two- dimensional turbulence: vortex clustering and merging,
and energy transfer to smaller scales



Direct cascade



Inverse cascade





Kolmogorov energy cascade

$$\frac{\text{kinetic energy } (\delta v_r)^2}{\text{time } r/\delta v_r} = \text{energy flux } \epsilon$$

$$\epsilon = \nu \langle |\nabla v|^2 \rangle \simeq v_{rms}^3/L$$

$$(\delta v_r)^3 \sim \epsilon r$$

$$S_3 = \langle (\delta v_r)^3 \rangle = -\frac{12\epsilon r}{d(d+2)}$$







FIG. 1. The experimental set-up.




We expect from turbulence
fragmentation, mixing and **loss of coherence**.

However,
an inverse turbulent cascade proceeds from small to large scales and brings some self-organization and eventually appearance of
a **coherent system-size condensate**.

Thin layer Condensation in two-dimensional turbulence

Temporal development of turbulence in a thin layer



$$E = E_0 e^{-\alpha t}$$







FIGURE 1. Time-averaged velocity field of the condensate in the square box of $L \approx 0.10$ m at different thicknesses of the bottom fluid (Fluorinert FC-77): (a) $\Delta h_b = 3$ mm, $\alpha = 0.3$ s $^{-1}$, (b) $\Delta h_b = 4$ mm, $\alpha = 0.15$ s $^{-1}$ and (c) $\Delta h_b = 5$ mm, $\alpha = 0.05$ s $^{-1}$. (d) Decay of the total kinetic energy for cases (a-c).

Mean subtraction recovers isotropic turbulence

1. Compute time-average velocity field (400 snapshots)

2. Subtract from 400 instantaneous velocity fields






Recover $\sim k^{-5/3}$ spectrum in the energy range

Kolmogorov law – linear $S_3(r)$ dependence in the “turbulence range”;

Kolmogorov constant $C \approx 7$

Universal profile of a coherent vortex



G. Boffetta, J. Laurie, I. Kolokolov,
V. Lebedev, GF, 2014




$$\mathbf{v} = (v_\phi, v_r) = (U + u, v) \text{ with } U(r), v(r, \phi, t), u(r, \phi, t)$$

$$\partial_t \mathbf{v} + \alpha \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = \mathbf{f} - \nabla p$$



$$r^{-2}\partial_r r^2\langle uv\rangle = -\alpha U$$


$$\epsilon = \frac{1}{r} \partial_r \left(r U \langle uv \rangle \right) + \alpha U^2$$

$$U^2 = 3\epsilon/\alpha$$



To understand atmosphere
one needs to move
from thin to **thick layers**






Video
Camera

square matrix of 30×30 permanent magnets (10 mm apart)

900 vortices (9 mm in diameter)

walls of the fluid cell ($0.3 \times 0.3 \text{ m}^2$)




Magnets

Electrodes


insertable square boundaries

$$L = (0.09 - 0.24) \text{ m}$$


$t = 5 \text{ s}$




$t = 5 \text{ s}$









$t = 20 \text{ s}$







Vertical shear suppresses
vertical vortices









single layer




double-layer





$$S_{3L} = \langle (\delta V_L)^3 \rangle$$

$$\varepsilon = -(2/3) S_{3L} / r$$



Moral

- A strong large-scale flow effectively suppresses fluctuations in the vertical velocity.
- The resulting flow is planar even at small scales yet it is three-dimensional as it depends strongly on the vertical coordinate.
- Turbulence in such flows transfers energy towards large scales.





Figure 1. Satellite image of hurricane Isabel, 12th September 2003 with the flight track of the WP-3D Orion aircraft (N43RF) overlaid. Stepped decent measurements of the boundary layer performed between outer rain bands.

Three- to two-dimensional turbulence transition in the hurricane boundary layer
D. Byrne and A. Zhang, 2014

A transition from 3d to 2d turbulence from in-situ aircraft measurements in the hurricane boundary layer



Third order structure function of horizontal velocities for different flight-leg heights in hurricane A) Isabel and B) Fabian.

These results represent the first measurement of the 2D upscale energy flux in the atmosphere and also the first to characterize the transition from 3D to 2D. It is shown that the large-scale parent vortex may gain energy directly from small-scales in tropical cyclones.

Summary

Inverse cascades seems to be scale invariant (and at least partially conformal invariant).

Condensation into a system-size coherent mode breaks symmetries of inverse cascades.

Condensates can enhance and suppress fluctuations in different systems. **Spectral condensates** of universal forms can coexist with turbulence.

Small-scale turbulence and large-scale vortex can conspire to provide for an inverse energy cascade.

Fluid Mechanics

The multi-disciplinary field of fluid mechanics is one of the most actively developing fields of physics, mathematics and engineering. In this book, the fundamental ideas of fluid mechanics are presented from a physics perspective.

Using examples taken from everyday life, from hydraulic jumps in a kitchen sink to Kelvin–Helmholtz instabilities in clouds, the book provides readers with a better understanding of the world around them. It teaches the art of fluid-mechanical estimates and shows how the ideas and methods developed to study the mechanics of fluids are used to analyse other systems with many degrees of freedom in statistical physics and field theory.

Aimed at undergraduate and graduate students, the book assumes no prior knowledge of the subject and only a basic understanding of vector calculus and analysis. It contains 32 exercises of varying difficulties, from simple estimates to elaborate calculations, with detailed solutions to help readers understand fluid mechanics.

Gregory Falkovich is a Professor in the Department of Physics of Complex Systems, Weizmann Institute of Science. He has researched in plasma, condensed matter, fluid mechanics, statistical and mathematical physics and cloud physics and meteorology, and has won several awards for his work.

Cover illustration: 'Sea Sky' © Rocksuzi, Dreamstime.com.

FALKOVICH
Fluid Mechanics

Fluid Mechanics

A Short Course for Physicists

GREGORY FALKOVICH

CAMBRIDGE
UNIVERSITY PRESS
www.cambridge.org



CAMBRIDGE

CAMBRIDGE

Weizmann Institute of Science, Israel

FEINBERG

GRADUATE SCHOOL

provides **housing** and
stipends for all students
and charges no tuition
fees

Application online
for **MSc** between Nov 1 and March 31
for **PhD** all year round

<https://www.weizmann.ac.il/feinberg>



Undergraduate Program

The **International Science Summer School** accepts a select number of outstanding ***undergraduate*** students each summer to participate in research projects in mathematics, physics, chemistry and the life sciences

Online application: Nov 1, 2015 -

Official results of the application:

March 15, 2016

Start of the program: June 2016

Duration of the program: 8 weeks



Kupcinet-Getz International Summer School

This is an enrichment program designed for undergraduate students from all over the world who are majoring in the sciences, and it is an opportunity to experience science by becoming part of a research group at the Weizmann Institute. Participation in the Kupcinet-Getz International Science School has had a significant impact on past participants of the program, reinforcing the students' interest in conducting research in their future careers.

As a student in this program, you will be associated with a laboratory or a theoretical research project under the supervision of an experienced scientist and, for a short period of time, will become a full-fledged member of the scientific team.

Stipends: NIS 500 (shekels) per week (up to **NIS 4,000~1100 \$** in total).

Accommodations: eight weeks of free rooming.

There are NO application, registration, visa or tuition fees.

Airfare or any other travel costs are not covered by the program. You are responsible for your own travel arrangements.